

AD-A118 122

TEXAS CHRISTIAN UNIV FORT WORTH DEPT OF CHEMISTRY
SYNTHESIS OF (SILYLAMINO)PHOSPHINES BY THE WILBURN METHOD.(U)
AUG 82 R H NEILSON, P WISIAN-NEILSON
TR-82-02

F/G 7/3

N00014-79-C-0632

NL

UNCLASSIFIED



END
DATE
FILMED
9 82
DTIC

AD A118122

OFFICE OF NAVAL RESEARCH

Contract N00014-79-C-0632

Task No. NR 053-720

TECHNICAL REPORT NO. 82-02

Synthesis of (Silylamino)phosphines By The

Wilburn Method

by

Robert H. Neilson and Patty Wisian-Neilson

Prepared for Publication

in

Inorganic Chemistry

Texas Christian University
Dept. of Chemistry
Fort Worth, TX 76129

August 1, 1982



Reproduction in whole or in part is
permitted for any purpose of the United States Government

This document has been approved for public
release and sale; its distribution is unlimited.

DTIC FILE COPY

82 08 09 065

unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 82-02	2. GOVT ACCESSION NO. AD-A118122	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Synthesis of (Silylamino)phosphines by the Wilburn Method		5. TYPE OF REPORT & PERIOD COVERED Technical Report 1981
		6. PERFORMING ORG. REPORT NUMBER TR-82-02
7. AUTHOR(s) R.H. Neilson and P. Wisian-Neilson		8. CONTRACT OR GRANT NUMBER(s) N00014-79-C00632
9. PERFORMING ORGANIZATION NAME AND ADDRESS Texas Christian University Fort Worth, TX 76129		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR-053-720
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research		12. REPORT DATE August 1, 1982
		13. NUMBER OF PAGES 6
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release: distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) Prepared for publication in <u>Inorganic Chemistry</u>		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Phosphine (Silylamino)phosphine		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Experimental details for the large scale (1-2 mol) synthesis of (silylamino)phosphines by the Wilburn method are described. In a "one-pot" procedure, treatment of PCl_3 with one equivalent of $LiN(SiMe_3)_2$ in Et_2O at $-78^\circ C$ followed by two equivalents of $RMgX$ at $0^\circ C$ affords 67-75% yields of $(Me_3Si)_2NPR_2$		

DD FORM 1473
1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014-6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

20. ABSTRACT

(R = Me, Et, CH_2SiMe_3). Similarly, $(\text{Me}_3\text{Si})_2\text{NP(Ph)R}$ (R = Me, Et) are obtained in 59-81% yields starting from PhPCl_2 . Physical and ^{31}P NMR data for these compounds are reported along with references to earlier papers describing their derivative chemistry.

Accession For	<input checked="checked" type="checkbox"/>
NTIS 03481	<input type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	<input type="checkbox"/>
By	
Distribution/	
Availability Codes	
Avail and/or	
Special	
Dist	
A	

DTIC
COPY
INSPECTED
2

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Contribution from the
Department of Chemistry
Texas Christian University
Fort Worth, Texas 76129

SYNTHESIS OF (SILYLAMINO)PHOSPHINES BY
THE WILBURN METHOD

Robert H. Neilson and Patty Wisian-Neilson

(Dedicated to the memory of Dr. James
C. Wilburn, 1953 - 1981)

Received _____

The importance of (silylamino)phosphines, e.g. $(\text{Me}_3\text{Si})_2\text{NPMe}_2$, as reagents for the synthesis of organo-substituted phosphazene polymers¹ and other novel organophosphorus compounds^{2,3} is now well established. Interest in these and related Si-N-P compounds as ligands in organometallic complexes⁴ is also developing and many other synthetic applications are likely to follow.

Most (silylamino)phosphines are prepared from commercially available reagents by a simple "one pot" synthesis first utilized by Wilburn.^{5,6} In earlier papers^{4a,6} we have described the preparation of a few specific compounds on relatively small scales (50-100 mmol). Because of the synthetic value of these reagents, however, we report here the complete details for the convenient, large-scale (ca. 1-2 mol) synthesis of (silylamino)phosphines via the Wilburn procedure.

Experimental Section

Materials and General Procedures. All reactions and other manipulations were carried out under an atmosphere of dry nitrogen. Ethyl ether was distilled from calcium hydride prior to use. The

following reagents were obtained from commercial sources: PCl_3 , PhPCl_2 , $(\text{Me}_3\text{Si})_2\text{NH}$, $n\text{-BuLi}$, MeMgBr , and EtMgBr . The silylmethyl Grignard $\text{Me}_3\text{SiCH}_2\text{MgCl}$ was prepared as needed according to the published procedure.⁷

Preparation of [Bis(trimethylsilyl)amino]dimethylphosphine (1).


A 5-L, 3-necked flask equipped with a paddle stirrer, a N_2 -flow, and a 500-mL addition funnel was charged with $(\text{Me}_3\text{Si})_2\text{NH}$ (1.0 mol, 209 mL) and Et_2O (ca. 1.0 L). One bottle of $n\text{-BuLi}$ (ca. 1.0 mol, 2.0 M in hexane) was transferred to the addition funnel under nitrogen pressure by means of a flexible, double-ended syringe needle⁸ and was added dropwise to the stirred silylamine solution at 0°C . The addition funnel was then washed with ca. 30 mL of ether and charged with PCl_3 (1.0 mol, 87.2 mL). After stirring the $\text{LiN}(\text{SiMe}_3)_2$ mixture at room temperature for ca. 90 min, it was cooled to -78°C and the PCl_3 was added dropwise. Upon completion of the addition, the -78°C bath was removed. During the warm-up period the orange colored solution gradually turned white as LiCl precipitated. After ca. one hour the mixture was cooled to 0°C and the addition funnel was again rinsed with Et_2O . Two bottles of MeMgBr (ca. 2.0 mol, 3.0 M in Et_2O) were transferred via the double-ended needle to the addition funnel and added dropwise to the stirred reaction mixture at 0°C over ca. 2 hours. The mixture was stirred for 3 hours and then allowed to stand over night at room temperature. Under a stream of N_2 , the supernatant solution was decanted into a large 1-necked flask. The solids were washed with two 500-mL portions of Et_2O and the washings were added to the first decantate. Most of the solvent was removed under reduced pressure and ca. 500 mL of hexane was added to facilitate precipitation of the remaining Grignard salts.

Filtration under N_2 followed by solvent removal left a viscous orange colored residue. Distillation through a short or semi-short path distilling head gave $(Me_3Si)_2NPMe_2$ ^{5,6} as a colorless liquid (166 g, 75% yield⁹, bp 55-60°C/4mm) of high purity based on 1H and ^{31}P NMR spectra.

Preparation of Other (Silylamino)phosphines. The same procedure using either commercial $EtMgBr$ or freshly prepared Me_3SiCH_2MgCl ⁷ affords the corresponding dialkylphosphines $(Me_3Si)_2NPR_2$, 2: $R=Et$ ¹⁰ and 3: $R=CH_2SiMe_3$.^{4a} Alternatively, if $PhPCl_2$ is used in place of PCl_3 and one equivalent of an alkyl Grignard is added in the last step, then the alkyl(phenyl)phosphines $(Me_3Si)_2NP(Ph)R$, 4: $R=Me$ ⁶ and 5: $R=Et$ ¹⁰, are obtained.

Results and Discussion

Treatment of a suspension of lithium bis(trimethylsilyl)amide in ether at -78°C with phosphorus trichloride followed by the addition of two equivalents of an alkyl Grignard reagent at 0°C affords the [bis(trimethylsilyl)amino]dialkylphosphines 1 - 3 (eq 1). Similarly, the use of dichlorophenylphosphine and one equivalent of Grignard reagent permits the convenient preparation of the corresponding alkyl(phenyl)phosphines 4, 5 (eq 2). With the exception of



some ^{31}P chemical shifts (Table I), the characterization of these compounds has generally been reported in other papers^{4a,6} describing their reactivity.

These reactions are easily carried out on large scales (ca. 1-2 mol) using unpurified commercial reagents. The products are obtained in relatively high yields (Table I) and good purity after a single vacuum distillation. Generally, the (silylamino)phosphines are colorless, foul-smelling liquids which are sensitive to oxidation and hydrolysis on exposure to the air. Nevertheless, their thermal stability is satisfactory and they can be stored indefinitely in tightly sealed containers.

This simple, "one-pot" synthesis is actually more general than is indicated here. For example, several different silylamines have been used^{5,6} and in favorable cases it is possible to obtain mono-alkylated products, e.g. $(\text{Me}_3\text{Si})_2\text{NP}(\text{Cl})\text{CH}_2\text{SiMe}_3$ ^{4a}. In more recent studies¹¹ involving very sterically congested systems, however, we find that reduced (P-H) or coupled (P-P) products are often obtained. Details of this work will be reported in subsequent papers.

Acknowledgment. This research is supported by the U.S. Army Research Office, the Office of Naval Research, and The Robert A. Welch Foundation.

References and Notes

- (a) Wisian-Neilson, P.; Neilson, R.H. J. Am. Chem. Soc. **1980**, 102, 2848.
 - (b) Neilson, R.H.; Wisian-Neilson, P. J. Macromol. Sci.-Chem. **1981**, A16, 425.
- Neilson, R.H.; Wisian-Neilson, P.; Morton, D.W.; O'Neal, H.R. ACS Symposium Series **1981**, 171, 239.
- (a) Neilson, R.H.; Goebel, D.W. J. Chem. Soc. Chem. Commun. **1979**, 769.
 - (b) Morton, D.W.; Neilson, R.H. Organometallics, **1982**, 1, 289.
- (a) Neilson, R.H. Inorg. Chem. **1981**, 20, 1679
 - (b) Thoma, R.J.; Ford, R.R.; Neilson, R.H. International Conference on Phosphorus Chemistry, Durham, N.C., June, **1981**. Abstract No. 247.
 - (c) Cowley, A.H.; Kemp, R.A.; Wilburn, J.C. Inorg. Chem. **1981**, 20, 4289.
 - (d) Scherer, O.J.; Jungmann, H. Angew. Chem. Int. Ed. Engl. **1979**, 18, 953.
 - (e) Keim, W.; Appel, R.; Storeck, A.; Kruger, C.; Goddard, R. Angew. Chem. Int. Ed. Engl. **1981**, 20, 116.
- Wilburn, J.C. Ph.D. Dissertation, Duke University, Durham, N.C., **1978**.
- Wilburn, J.C.; Neilson, R.H. Inorg. Chem. **1979**, 18, 347.
- Andersen, R.A.; Wilkinson, G. Inorg. Synth. **1979**, 19, 262.
- Shriver, D.F. "The Manipulation of Air-Sensitive Compounds," McGraw-Hill, New York, **1969**; p. 156.
- Yields up to 86% have been obtained by more careful and repeated washing of the solids.
- ¹H NMR data: 2, δ 0.34 (Me₃Si, J_{PH} = 0.6 Hz), δ 1.14 (CH₃, J_{PH} = 16.0 Hz, J_{HH} = 7.5 Hz), δ 1.4-2.0 (CH₂, multiplet). 5, δ 0.25 (Me₃Si, J_{PH} = 0.6 Hz), δ 1.33 (CH₃, J_{PH} = 18.0 Hz, J_{HH} = 7.2 Hz), δ 1.9-2.2 (CH₂, multiplet), δ 7.1-7.4 (Ph, multiplet)
- manuscripts in preparation.

Table I. Physical and ^{31}P NMR Data for Some
(Silylamino)phosphines, $(\text{Me}_3\text{Si})_2\text{NPRR}'$

Compd	R	R'	bp (mm)	Yield (%)	$\delta(^{31}\text{P})^a$
1	Me	Me	55-60 $^\circ$ (4)	75	31.7
2	Et	Et	68-69 $^\circ$ (1.6)	71	57.2
3	CH_2SiMe_3	CH_2SiMe_3	80-85 $^\circ$ (0.3)	67	42.9
4	Me	Ph	90-95 $^\circ$ (0.8)	81	37.6
5	Et	Ph	85-90 $^\circ$ (0.1)	59	50.9

^aChemical shifts downfield from external H_3PO_4 , measured in CDCl_3 solution.

Abstract

Experimental details for the large scale (1-2 mol) synthesis of (silylamino)phosphines by the Wilburn method are described. In a "one-pot" procedure, treatment of PCl_3 with one equivalent of $\text{LiN}(\text{SiMe}_3)_2$ in Et_2O at -78°C followed by two equivalents of RMgX at 0°C affords 67-75% yields of $(\text{Me}_3\text{Si})_2\text{NPR}_2$ ($\text{R} = \text{Me}, \text{Et}, \text{CH}_2\text{SiMe}_3$). Similarly, $(\text{Me}_3\text{Si})_2\text{NP}(\text{Ph})\text{R}$ ($\text{R} = \text{Me}, \text{Et}$) are obtained in 59-81% yields starting from PhPCl_2 . Physical and ^{31}P NMR data for these compounds are reported along with references to earlier papers describing their derivative chemistry.

